

The Tap – Midazolam was Used to Prematurely End the Lives of Thousands who You were Told had Died of Covid 19 and We can Prove It

 robscholtemuseum.nl/the-tap-midazolam-was-used-to-prematurely-end-the-lives-of-thousands-who-you-were-told-had-died-of-covid-19-and-we-can-prove-it-heres-the-evidence

29 augustus 2021

Farmacia (1) (foto Cercafarmaco)

Farmacia (2) (foto Il Viaggiatore)


Farmacia (3) (foto cantabriatotal.com)

Farmacia (4) (foto comuneterelle.it)

Farmacia (5) (foto Farmacia Zorzetto Paolo Ca' Savio Venezi)

Farmacia (6) (foto Pharmacia Ferrari)

Farmacia (7) (foto Brevetto Farmaco di Alessandria)



Midazolam 2D Structure (foto Mol-Instincts)


Ball & Stick Model Image of Optimized Midazolam Structur (foto Mol-Instincts)

Midozolam (foto Mol-Instincts)

van der Waals Surface Animated Gif of Optimized Midazolam Structure (foto Mol-Instincts)

Midazolam 3D (foto Mol-Instincts)





Midazolam Injectable (foto telegra.ph)

An advertisement for Flormidal Midazolam. It features a woman with blonde hair sleeping peacefully on a white pillow. In the foreground, there are several blister packs of Flormidal 15 mg tablets. A cloud-like graphic contains the text "15 mg". A large, stylized watermark "www.pharmacy-weightloss.net" is overlaid diagonally across the image. The product name "Flormidal Midazolam" is written in a large, bold, red font at the top left of the advertisement area.






Midazolam





Una Reaccion Paradojica di Midozolam (foto Imgflip)





The Real First Wave Midazolam (foto brandnewtube.com)

But what if we could prove to you that you've given up fifteen months and counting of your life due to a lie? But not just any lie, a lie that has involved prematurely ending the lives of thousands upon thousands of people, who you were told died of Covid-19. A lie that has involved committing one of the greatest crimes against humanity in living memory. A lie that has required three things – fear, your compliance, and a drug known as Midazolam... Authorities state that Covid-19 is an infectious disease caused by a new coronavirus dubbed SARS-CoV-2. The World Health Organisation (WHO) tell us that "most people infected with the COVID-19 virus will experience mild to moderate respiratory illness and recover without requiring special treatment". However they state that "older people, and those with underlying medical problems like cardiovascular disease, diabetes, chronic respiratory disease, and cancer are more likely to develop serious illness". We're told that serious illness in Covid-19 presents pneumonia and accompanying respiratory insufficiency. Therefore typical symptoms include breathlessness, cough, weakness and fever. We're also told that people who suffer deteriorating respiratory failure and who do not receive intensive care, develop acute respiratory distress syndrome with severe breathlessness.

Recommendations for Symptom Control of Patients with COVID-19

Issue


COVID-19 is an acute disease with a clinical presentation of pneumonia and accompanying respiratory insufficiency. Thus, typical symptoms are dyspnoea (breathlessness), cough, weakness and fever. Other symptoms such as anxiety, panic, restlessness and delirium have been reported. Patients with rapidly deteriorating respiratory failure and who do not receive intensive care, develop acute respiratory distress syndrome (ARDS) with severe breathlessness, anxiety and panic, requiring rapid intervention for symptom control.

Pneumonia is an inflammation of one or both lungs, usually caused by an infection. It causes the alveoli (air sacs) inside the lungs to fill with fluid, making it harder for them to work properly. The body sends white blood cells to fight the infection, and while this helps kill the germs it can also make it harder for the lungs to pass oxygen into the bloodstream.

Pneumonia is not a new condition that has appeared due to Covid-19. In 2019 alone, the year prior to the alleged emergence of Covid-19, [272,000 people were admitted to hospital](#) with pneumonia. According to the British Lung Foundation [in 2012, 345 people per every 100,000](#) had one or more episodes of pneumonia. This equates to around 225,000 people suffering pneumoni at least once.

The British Lung Foundation also show us that the majority of cases of pneumonia occur in those who are aged 81 and over. For instance in 2012 1,838 people in every 100,000 people over the age of 81 developed pneumonia, this equates to around 60,000 people over the age of 81 in today's numbers based on there being around [3.2 million people over the age of 80](#) in the UK.

Number of people per 100,000 who developed pneumonia, by age group, 2004–12




Source [British Lung Foundation](#)

In March 2020 the British people were told that they must “stay at home” in order to “protect the NHS” and “save lives”. They were also told that the authorities needed just “three weeks to flatten the curve”.


They also tell us that In 2012 there were 28,592 deaths from pneumonia, which equates to 5.1% of all deaths that occurred that year.

UK deaths from pneumonia compared with other lung diseases, 2012



So as you can see deaths due to pneumonia have not just suddenly started to happen because of the alleged emergence of a new disease called Covid-19, we've been living with them all of our lives, they just haven't been thrust in front of you 24/7 on the television, or the front page of every newspaper like they have with alleged Covid-19 deaths.

But for us to prove to you that you given up fifteen months of your life due to a lie that involved prematurely ending the lives of thousands upon thousands of people we first need you to understand which age group has been most affected by Covid-19 according to official statistics.



The above graph is a heat map showing deaths within 28 days of a positive test for SARS-CoV-2 by date of death and age of the person. This data can be seen at the UK Gov's coronavirus dashboard [here](#). What's pretty clear from this data is that the most alleged Covid deaths have occurred in people aged 90+. The next age group with the most deaths being 85 – 89, then 80 – 84 and so on and so on. There's a general decrease in the number of deaths up to about the 65-69 age group but then we see a dramatic fall to pretty much zero in anyone aged under about the age of 60.

This heat map shows that there have generally been no more than 9 deaths in a single day of anyone aged between 60 – 64. In the 65-69 year old group there have been no more than 20 deaths a day. In the 70-74 year old group no more than 27 deaths in a day. In the 75-79 group no more than 48 deaths in a day, at it's highest. It isn't until we get to the 85-89 year old group that we start to see a large increase in the number of alleged Covid deaths. 179 deaths in a day at its highest. Then we have the 90+ age group which has seen no more than 379 deaths in a single day at it's highest.

So what we're seeing here is that is a negligible amount of "Covid" deaths in anyone under the age of 60. But we're really not seeing very many "Covid" deaths in anybody aged between 60 and 80. What we are seeing is a much higher amount of "Covid" deaths in people aged over 85. But what's so strange about that?

Well nothing when we consider the average life expectancy in the UK is 81 years of age. Plus the fact this is also in line with what we have seen in cases / deaths due to pneumonia in previous years. Don't forget serious illness in Covid-19 presents **pneumonia** and accompanying respiratory insufficiency.

Which begs the question of why the entire nation has had to stay at home, social distance, wear a mask, wash their hands, and live under dictatorial tyranny for fifteen months because people who have lived longer than the average life expectancy in the UK have died or are dying? Dying of pneumonia from which we see tens of thousands of deaths every year/

We cannot deny that 2020 did see excess deaths, and you would believe this is due to the hospitals being overwhelmed? Except they weren't.

[NHS data](#) shows us that during the height of the "first wave" between April and June 2020 there were 58,005 beds occupied which equated to 62% occupancy. This is 30% down on the same time frame in the previous year.

Title:	Average daily number of available and occupied beds open overnight by sector			
Summary:	KH03 is the collection of data to monitor available and occupied beds open overnight that are consultant led.			
Period:	April to June 2020			
Source:	NHS England: SDCS data collection - KH03			
Basis:	Provider			
Published:	20th August 2020			
Revised:	19th November 2020			
Status:	Public			
Contact:	Paul Steele - england.nhsdata@nhs.net			

Occupied	
General & Acute	Learning Disabilities
58,005	602

% Occupied	
General & Acute	Learning Disabilities
62.7%	70.6%

- In 2017, April-June there were on average a total of 91,724 beds occupied which equated to 89.1% occupancy.
- In 2018, April-June there were on average a total of 91,056 beds occupied which equated to 89.8% occupancy.
- In 2019, April-June there were on average a total of **91,730** beds occupied which equated to **90.3%** occupancy.
- **In 2020, April-June there were on average a total of 58,005 beds occupied which equated to 62% occupancy.**

It also shows us that [A&E attendance](#) during the height of the first wave was 57% down on the previous year.

Summary: A&E attendances, performance and emergency admissions

Period: April 2020

Source: SDCS data collection - MSitAE

Basis: Provider

Published: 14th May 2020

Revised:

Status: Published

Contact: Chris Evison - England.nhsdata@nhs.net

A&E attendances			
Type 1 Departments - Major A&E	Type 2 Departments - Single Specialty	Type 3 Departments - Other A&E/ Minor Injury Unit	Total attendances
689,720	19,726	207,135	916,581

Provider Level Data


- 2018 – April – 1,984,369 attended A&E
- 2019 – April – **2,112,165** attended A&E
- **2020 – April – 916,581 attended A&E**

Which begs the question of what exactly were we protecting the NHS from? It seems to have had a holiday.

But there were 41,627 more deaths than the five year average up to the 1st May 2020, and the vast majority of these occurred in April. An April which saw A&E attendance down 57% compared to the previous year and bed occupancy down 30% compared to the previous year. 33,408 of these excess deaths mentioned Covid-19 on the death certificate, the vast majority of which occurred in those over the age of 85,

Figure 1: The total number of deaths (all cause) and the number of deaths involving COVID-19 decreased for the second week running

Number of deaths registered by week, England and Wales, 28 December 2019 to 1 May 2020



Source: Office for National Statistics

However [data taken from the Office for National Statistics \(ONS\)](#) shows us that during April 2020 there were 26,541 deaths that occurred in care homes, an increase of 17,850 on the five-year average. This is half the amount of alleged Covid-19 deaths during the same period.

	Month of occurrence	Five-year average (2015-2019)		2020	
		Home	Care home	Home	Care home
England	January	11,503.8	12,059.4	11,319	10,831
England	February	9,897.0	10,022.2	10,348	9,522
England	March	10,316.6	10,166.6	12,766	11,827
England	April	9,384.6	8,691.0	16,909	26,541
England	May	9,491.4	8,401.0	13,308	13,953
England	June	8,917.0	7,809.2	12,096	7,971
England	July	9,191.0	8,089.0	11,832	7,557
England	August	9,093.6	8,169.0	11,663	7,838
England	September	8,877.2	8,099.4	11,130	7,775
England	October	9,779.0	9,239.2	12,500	9,119
England	November	10,002.4	9,372.2	12,365	9,676
England	December	11,135.0	10,830.0	12,657	10,335

Why did so many people die in care homes when hospitals were far from overwhelmed? Surely if they have developed serious complications due to Covid-19 they would require urgent medical attention and hospital treatment?

Because don't forget we're told that serious illness in Covid-19 presents **pneumonia** and accompanying respiratory insufficiency. Therefore typical symptoms include **breathlessness**, cough, weakness and fever. We're also told that people who suffer deteriorating **respiratory failure** and who do not receive **intensive care**, develop acute **respiratory distress** syndrome with **severe breathlessness**.

Why were these people in care homes and not in hospital?

They were in care homes because Matt Hancock gave the order to put them there...

1. Summary

- 1.1 This document sets out the Hospital Discharge Service Requirements for all NHS trusts, community interest companies and private care providers of acute, community beds and community health services and social care staff in England, who must adhere to this from Thursday 19th March 2020. It also sets out requirements around discharge for health and social care commissioners (including Clinical Commissioning Groups and local authorities).
- 1.2 Unless required to be in hospital (see Annex B), patients must not remain in an NHS bed.
- 1.3 Based on these criteria, acute and community hospitals must discharge all patients as soon as they are clinically safe to do so. Transfer from the ward should happen within one hour of that decision being made to a designated discharge area. Discharge from hospital should happen as soon after that as possible, normally within 2 hours.
- 1.4 Implementing these Service Requirements is expected to free up to at least 15,000 beds by Friday 27th March 2020, with discharge flows maintained after that. Acute and community hospitals must keep a list of all those suitable for discharge and report on the number and percentage of patients on the list who have left the hospital and the number of delayed discharges through the daily situation report.

On the 19th March [a directive was sent out to the NHS](#) which required them to discharge all patients who they deemed to not require a hospital bed. They declared that transfers from the ward must happen within one hour of that decision being made to a designated discharge area, and that discharge from hospital should happen within 2 hours. NHS trusts were told that "they must adhere" to the new directive.

This was done to allegedly free up beds, of which they estimated would amount to an extra 15,000 free beds within just one week of the directive being implemented.

It freed up so many beds that bed occupancy during April – June 2020 was 30% down on the previous year. Why on earth would these people already be in a hospital bed if they did not need to be? You attend hospital because you require medical treatment, not because you want a lie down and a good nights sleep.

This directive meant that people who required medical treatment and attention were discharged into Care homes in the thousands.

But Matt Hancock's abandonment of the elderly and vulnerable didn't end there. Whilst the NHS was busy discharging patients who required medical treatment into care homes under his directive, Matt Hancock and the Department of Health were busy trying to source them all a certain drug known as **midazolam**.



Midazolam is a commonly used drug in palliative care and is considered one of the four essential drugs needed for the promotion of quality care in dying patients in the United Kingdom. Think of it as diazepam on steroids.

Midazolam is also a drug that has been used in executions by lethal injection in the USA, combined with two other drugs. Midazolam acts as a sedative to make the prisoner unconscious. The other drugs then stop the lungs and heart working. However it has been the source of controversy as several prisoners took a long while to die and appeared to be in pain when midazolam was used.

Midazolam can also cause serious or life-threatening breathing problems such as shallow, slowed, or temporarily stopped breathing that may lead to permanent brain injury or death.

UK regulators state that you should only receive midazolam in a hospital or doctor's office that has the equipment that is needed to monitor your heart and lungs and to provide life-saving medical treatment quickly if your breathing slows or stops.

A doctor or nurse should watch you closely after you receive this medication to make sure that you are breathing properly because midazolam induces significant depression of respiration. Your doctor should also be made aware if you have a severe infection or if you have or have ever had any lung, airway, or breathing problems or heart disease.

Midazolam is also used before medical procedures and surgery to cause drowsiness, relieve anxiety, and prevent any memory of the event. It is also sometimes given as part of the anesthesia during surgery to produce a loss of consciousness.

Midazolam is also used to cause a state of decreased consciousness in seriously ill people in intensive care units who are breathing with the help of a machine.

Midazolam should be used with extreme caution in patients who have chronic renal failure, impaired hepatic function, or impaired cardiac function. It should also be used with extreme caution in obese patients, or elderly patients.

What are some of the most important points you should take from this?

- Midazolam induces significant depression of respiration
- UK regulators insist midazolam should only be administered in a hospital or doctor's office under the supervision of a doctor or nurse to monitor the breathing of the patient in order to provide life saving treatment to the patient if breathing slows or stops.
- Midazolam should be used with extreme caution in elderly patients

Serious illness in Covid-19 presents pneumonia and accompanying respiratory insufficiency. Therefore typical symptoms include breathlessness, cough, weakness and fever. We're also told that people who suffer deteriorating respiratory failure and who do not receive intensive care, develop acute respiratory distress syndrome with severe breathlessness.

Midazolam induces significant depression of respiration.

Knowing that would you use midazolam to treat people who were suffering pneumonia and respiratory insufficiency allegedly due to Covid-19?

WARNINGS

Personnel and Equipment for Monitoring and Resuscitation

Adults and Pediatrics: Intravenous midazolam hydrochloride has been associated with respiratory depression and respiratory arrest, especially when used for sedation in noncritical care settings. In some cases, where this was not recognized promptly and treated effectively, death or hypoxic encephalopathy has resulted. Intravenous midazolam hydrochloride should be used only in hospital or ambulatory care settings, including physicians' and dental offices, that provide for continuous monitoring of respiratory and cardiac function, e.g., pulse oximetry. Immediate availability of resuscitative drugs and age- and size-appropriate equipment for bag/valve/mask ventilation and intubation, and personnel trained in their use and skilled in airway management should be assured (see [WARNINGS](#)). For deeply sedated pediatric patients, a dedicated individual, other than the practitioner performing the procedure, should monitor the patient throughout the procedure.

Risks From Concomitant Use With Opioids

Concomitant use of benzodiazepines and opioids may result in profound sedation, respiratory depression, coma, and death. Monitor patients for respiratory depression and sedation (see [WARNINGS](#) and [PRECAUTIONS, DRUG INTERACTIONS](#)).

[Source – US National Library of Medicine](#)

Well Matt Hancock and friends certainly seem to think so as you can see in the following video... https://www.youtube.com/embed/s-dqiRr-o5A?version=3&rel=1&showsearch=0&showinfo=1&iv_load_policy=1&fs=1&hl=en-US&autopause=2&wmode=transparent

Q377 Dr Evans: A good death needs three things: equipment, medication and the staff to administer it. On equipment, do you have enough syringe drivers in the NHS to deliver medications to keep people comfortable when they are passing away?

Matt Hancock: Yes, we have. A challenge was raised on that about eight days ago—it was not as big a challenge as was made public, and we have resolved it. Yes; right now we have enough.

Q378 Dr Evans: The syringe drivers are used to deliver medications such as midazolam and morphine. Do you have any precautions in place to ensure that we have enough of those medications?

Matt Hancock: Yes. We have a big project to make sure that the global supply chains for those sorts of medications, as well as the ITU medications that I spoke about earlier, are clear. In fact, those medicines are made in a relatively small number of factories around the world, so it is a delicate supply chain and we are in contact with the whole supply chain.

[Source](#)

The above exchange took place in a parliamentary committee meeting on the 17th April 2020 between Matt Hancock and Dr Evans, who is a fellow Conservative MP.

The following is an extract from [an article](#) which confirms the United Kingdom purchased two years worth of Midazolam in March 2020 and were looking to purchase much more –

Supplies of the sedative midazolam have been diverted from France as a “precaution” to mitigate potential shortages in the NHS caused by COVID-19, the Department of Health and Social Care (DHSC) has told The Pharmaceutical Journal.

A spokesperson from Accord Healthcare, one of five manufacturers of the drug, told The Pharmaceutical Journal that it had to gain regulatory approval to sell French-labelled supplies of midazolam injection to the NHS, after having already sold two years’ worth of stock to UK wholesalers “at the request of the NHS” in March 2020.

The DHSC said the request for extra stock was part of “national efforts to respond to the coronavirus outbreak”, which included precautions “to reduce the likelihood of future shortages”.

Why on earth would the United Kingdom need to purchase two years worth of Midazolam, a drug associated with respiratory suppression and respiratory arrest, to treat a disease that causes respiratory suppression and respiratory arrest?

Individualization of Dosage

Midazolam hydrochloride must never be used without individualization of dosage. The initial intravenous dose for sedation in adult patients may be as little as 1 mg, but should not exceed 2.5 mg in a normal healthy adult. Lower doses are necessary for older (over 60 years) or debilitated patients and in patients receiving concomitant narcotics or other central nervous system (CNS) depressants. The initial dose and all subsequent doses should always be titrated slowly; administer over at least 2 minutes and allow an additional 2 or more minutes to fully evaluate the sedative effect. The use of the 1 mg/mL formulation or dilution of the 1 mg/mL or 5 mg/mL formulation is recommended to facilitate slower injection. Doses of sedative medications in pediatric patients must be calculated on a mg/kg basis, and initial doses and all subsequent doses should always be titrated slowly. The initial pediatric dose of midazolam for sedation/anxiolysis/amnesia is age, procedure, and route dependent (see [DOSAGE AND ADMINISTRATION](#) for complete dosing information).

[Source – US National Library of Medicine](#)

This [document](#) produced by the NHS states that Midazolam should be used for comfort at end of life care due to Covid-19 to ease fear, anxiety and agitation. [Source](#)

University Hospitals Birmingham 
NHS Foundation Trust

Benzodiazepines = FIRST LINE for anxiety, fear and agitation

Midazolam – suggest start with **low doses** for patients naïve to this drug but be prepared, if response is poor or short lived and anxiety is severe, to **escalate dosing sharply if required**.

- **Generally:** Start with 2.5 mg SC or IV
- If patient is **particularly frail**: use 1.25mg
- If **extremely distressed** or show **tolerance** to this group of drugs: may require higher doses e.g. 5 -10 mg

If ward areas **cannot access midazolam** then lorazepam can be used as a substitute – generally **2.5 mg of midazolam can be regarded as 'equivalent' to 500 mcg of injectable lorazepam**.
Seek advice.

For patients not responding to midazolam – this might be because doses have been too low or not frequent enough. Some patients might need much higher doses than normal. **Seek advice** if needed.

This [NHS document](#) states that midazolam should be used for sedation prior to the patient requiring mechanical ventilation, something we know has been required in hospitals for people who have developed severe pneumonia, of which we are told is due to Covid-19. However it also states that midazolam should only be used if 1st line and 2nd line drugs do not provide adequate sedation, but does include the caveat that midazolam alone can be added to 1st line drugs to reduce Propofol infusion rates. [Source](#)

3rd Line

Morphine 0.04-0.2 mg/kg/hour

Midazolam 0.04-0.2mg/kg/hour

This regimen may be used if 1st and 2nd line drugs fail to provide adequate sedation without unacceptable cardiovascular depression or Midazolam alone can be added to the first line drugs to reduce Propofol infusion rates.

This [NHS document](#) states that midazolam should be used for sedation prior to having a operation. [Source](#)

Drugs preferred for sedation

The preferred drugs to be used for sedation are midazolam and fentanyl. These are both short acting drugs with an onset of 1-2 minutes and a peak effect of 3-5 minutes.

The analgesic effects of fentanyl last about 30-60 minutes therefore occasionally morphine is needed for a more prolonged analgesic effect after the procedure.

Morphine has a longer onset with a peak intravenous effect of 20 minutes, therefore the patients should be observed for at least 10-15 minutes before the midazolam is administered. See appendix 2 for further information about dosing, side effects and cautions. Propofol should not be used by non-anaesthetists for sedation.

The same document also provides confirmation that midazolam has the potential to impair the respiration system, particularly in the presence of disease or old age. It clearly states that dosage should be kept to a minimum and should be within the manufacturer's guidelines.

The drugs used for IV conscious sedation are benzodiazepines and opioids. **These drugs depress the central nervous system and have the potential to impair respiration or circulation or both,** particularly in the presence of disease or old age.³ When used together there is a synergistic effect which can increase the risk of cardiorespiratory events. Dosage of benzodiazepines and opiates should be kept to a minimum to achieve sedation and should be within the manufacturer's guidelines.⁴

The document also provides a helpful table confirming dosage of midazolam for the elderly or unwell should be no more than 0.5mg – 1 mg, side effects include cardiorespiratory depression and the drug should be used with caution in those suffering respiratory disease.

Appendix 2

Table of Preferred Drugs - list including safe dosages

Sedative	Dose	Onset	Side effects	Cautions
Midazolam	<u>Healthy adult</u> : 1-2mg bolus. Titrate further small boluses with at least 2 minutes between doses. Usually max 5mg required. <u>Elderly or unwell</u> : 0.5-1mg bolus. Titrate further small boluses with at least 2 minutes between doses. Often no more than 2 mg required.	3-5 minutes for peak effect. Half life 1.5-3.5 hours.	CARDIORESPIRATORY DEPRESSION especially associated with opioids. Gastro-intestinal disturbances, anaphylaxis, drowsiness, confusion, ataxia, amnesia, headache, paradoxical excitement and aggression (especially in children and elderly), dysarthria; injection-site reactions. For complete list see BNF	Cardiac disease; hepatic impairment; renal impairment; (increases plasma half life x2-2.5) respiratory disease; myasthenia gravis; history of drug or alcohol abuse; risk of severe hypotension in hypovolaemia, vasoconstriction, hypothermia; pregnancy and breast-feeding

This article confirms that over 2 million operations were cancelled at the end of March 2020 to free up beds for at least three months for “coronavirus” patients. – [Source](#)

More than 2m operations cancelled as NHS fights Covid-19

Backlog of procedures in England could cost NHS £3bn to work through after crisis

- [Coronavirus - latest updates](#)
- [See all our coronavirus coverage](#)

Ian Sample *Science editor*

🐦 @iansample

Sun 26 Apr 2020 16.43 BST



Can you see the contradictions here? A policy that has been in place prior to the alleged emergence of Covid-19 clearly states that midazolam can be used for sedation, however dosage should be reduced to 0.5mg in the elderly or unwell due to possible side effects which include cardiorespiratory depression, and extreme caution should be used in administering midazolam to patients suffering respiratory disease.

However a policy created for treating patients allegedly suffering anxiety due to Covid-19, which we're told is a respiratory disease, clearly states to treat said patient with a starting dose of 2.5mg of Midazolam, or 1.25mg if the patient is "particularly frail", but to bump this up to 5 – 10mg if the patient is "extremely distressed". Even the starting dose for the particularly frail is 0.25mg higher than the maximum recommended to administer to the elderly or unwell in sedation guidelines.

Who is responsible for making this decision and publishing these guidelines? And why is nobody holding them to account?

Two years worth of Midazolam was purchased in March 2020, however at the same time operations were cancelled for a minimum of three months, therefore Midazolam was not required for use in sedation prior to operations. Guidelines published prior to the alleged pandemic clearly state that Midazolam should be used in extremely low doses in the elderly or unwell, and should be used with extreme caution in those suffering respiratory disease due to side effects which include respiratory depression. We're told Covid-19 is a respiratory disease and complications present pneumonia and severe respiratory distress. Therefore considering all of this the purchase of two years worth of Midazolam seems to be an awful waste of money, doesn't it? As there doesn't seem to be much they could possibly use it for within the guidelines

Well we can confirm it was definitely used as we have seen the prescription data.

But we'd just like to remind you of the [important warning applied to Midazolam](#) courtesy of the US National Library of Medicine –

Midazolam injection may cause serious or life-threatening breathing problems such as shallow, slowed, or temporarily stopped breathing that may lead to permanent brain injury or death. You should only receive this medication in a hospital or doctor's office that has the equipment that is needed to monitor your heart and lungs and to provide life-saving medical treatment quickly if your breathing slows or stops. Your doctor or nurse will watch you closely after you receive this medication to make sure that you are breathing properly.

So can Matt Hancock explain why during April 2020 out of hospital prescribing for Midazolam was twice the amount seen in 2019?

Midazolam Hydrochloride (1501041T0)


Part of chapter [15 Anaesthesia](#), section [15.1 General anaesthesia](#), paragraph [15.1.4 Sedative and analgesic peri-operative drgs](#)

High-level prescribing trends for Midazolam Hydrochloride (BNF code 1501041T0) across all GP practices in NHS England for the last five years. You can see [which CCGs prescribe most of this chemical](#) relative to its class, or learn more [about this site](#).


[View all matching dm+d items.](#)

Trends

Items Spending




According to [official data](#) in April 2019 up to 21,977 prescriptions for Midazolam were issued, containing 171,952 items, the vast majority being Midazolam Hydrochloride. However in April 2020 45,033 prescriptions for Midazolam were issued, containing 333,229 items, the vast majority being Midazolam Hydrochloride. That is a 104.91% increase in the number of prescriptions issued for Midazolam and a 93.85% increase in the number of items they contained. But these weren't issued in hospitals, they were issued by GP practices which can only mean one thing, they were issued for end of life care.



The above is a graph displayed on the UK Government website displaying deaths within 28 days of a positive test result for Covid-19 by date of death.

The following graph has been created using data on the amount of Midazolam solution produced each month from January 20219 through to March 2021.



Can you spot the difference? We couldn't either because there isn't one.

The spikes in production of Midazolam solution match the spikes of alleged Covid deaths within 28 days of a positive test.

April 2020 – huge surge in Midazolam prescriptions out of hospital and huge surge in production of Midazolam solution.

April 2020 – huge surge in alleged Covid deaths.

January 2021 – huge surge in production of Midazolam solution.

January 2021 – huge surge in alleged Covid deaths.

We're told that serious illness in Covid-19 presents pneumonia and accompanying respiratory insufficiency. Therefore typical symptoms include breathlessness, cough, weakness and fever. We're also told that people who suffer deteriorating respiratory failure and who do not receive intensive care, develop acute respiratory distress syndrome with severe breathlessness.

Midazolam Hydrochloride is associated with respiratory depression and respiratory arrest, especially when used for sedation in noncritical care settings. In some cases, where this was not recognized promptly and treated effectively, death or hypoxic encephalopathy has resulted. Intravenous midazolam hydrochloride should be used only in hospital or ambulatory care settings.

NHS policy prior to the emergence of Covid-19 states –

Dosage should be reduced to 0.5mg in the elderly or unwell due to possible side effects which include cardiorespiratory depression, and extreme caution should be used in administering midazolam to patients suffering respiratory disease.

NHS policy after the emergence of Covid-19, an alleged respiratory disease states

Benzodiazepines = FIRST LINE for anxiety, fear and agitation

Midazolam – suggest start with **low doses** for patients naïve to this drug but be prepared, if response is poor or short lived and anxiety is severe, to **escalate dosing sharply if required**.

- **Generally:** Start with 2.5 mg SC or IV
- If patient is **particularly frail**: use 1.25mg
- If **extremely distressed** or show **tolerance** to this group of drugs: may require higher doses e.g. 5 -10 mg

If ward areas **cannot access midazolam** then lorazepam can be used as a substitute – generally 2.5 mg of midazolam can be regarded as ‘equivalent’ to 500 mcg of injectable lorazepam.
Seek advice.

For patients not responding to midazolam – this might be because doses have been too low or not frequent enough. Some patients might need much higher doses than normal. **Seek advice** if needed.

Hospitals beds in April 2020 30% were down compared to the previous year.

A&E attendance was 57% down in April 2020 compared to the previous year.

Care home deaths were 205% up in April 2020 compared to April 2019.

The vast majority of alleged Covid deaths are people over the age of 85.

Can you not see a strong correlation here between the over prescribing of Midazolam and the seemingly premature ending of life, with the associated deaths being put down as Covid-19?

Did you really believe there's a virus so clever that it knows to kill people who are disabled? Just look at the ONS statistics. Three in every five alleged Covid-19 deaths occurred in those who suffered learning difficulties and disabilities ([see here](#)).

Disabled people made up 6 in 10 (59.5%) of all deaths involving the coronavirus (COVID-19) for the period to 20 November 2020 (30,296 of 50,888 deaths). For comparison, disabled people made up 17.2% of the study population, therefore suggesting that disabled people have been disproportionately impacted by the COVID-19 pandemic. See the [datasets](#) for detailed breakdowns of counts of deaths.

In relation to deaths of people with learning difficulties the ONS said – *‘the largest effect was associated with **living in a care home** or other communal establishment.*’

Having a learning difficulty and being in care doesn’t mean you are more likely to die of Covid-19. What it means is that you are much more likely to have a DNR order placed on you without informing yourself or your family, which Carers / NHS staff then use as permission to put you on end of life care, which involves the administration of Midazolam.

We know this happened because an [Amnesty report](#) and [CQC report](#) said so.

DENIAL OF ACCESS TO HOSPITALS AND OTHER MEDICAL SERVICES

Amnesty International has received multiple reports of care home residents’ right to NHS services, including access to general medical services (GMS) and hospital admission, being denied during the pandemic, violating their right to health and potentially their right to life, as well as their right to non-discrimination. Care homes managers have pointed out that such reluctance or refusal to admit older care home residents to hospital could not be explained by need, as hospital bed capacity was never reached.¹¹⁰

The Amnesty Report states that

‘Care home managers and staff and relatives of care home residents in different parts of the country told Amnesty International how, in their experience, sending residents to hospital was discouraged or outright refused by hospitals, ambulance teams, and GPs. A manager in Yorkshire said: “We were heavily discouraged from sending residents to hospital. We talked about it in meetings; we were all aware of this.”’

‘Another manager in Hampshire recalled:

There wasn’t much option to send people to hospital. We managed to send one patient to hospital because the nurse was very firm and insisted that the lady was too uncomfortable and we could not do any more to make her more comfortable but the hospital could. In hospital the lady tested COVID positive and was treated and survived and came back. She is 92 and in great shape.

She explained that:

There was a presumption that people in care homes would all die if they got COVID, which is wrong. It shows how little the government knows about the reality of care homes.’

The nurse from the GP surgery rang me up to say they decided mum is DNR. I asked why and she said “we did this across the home”, and I said “no, this should be done on individual cases and I don’t agree to it”. So I had it taken off ... She also said that they would not take mum to hospital and again I said that is something that would have to be decided if and when need arose on the basis of the situation at the time. They had asked mum about the DNR and she had agreed to it but then I spoke to mum and she had not really understood the issue.

‘The son of one care home resident who passed away in Cumbria said that sending his father to hospital had not even been considered:

From day one, the care home was categoric it was probably COVID and he would die of it and he would not be taken to hospital. He only had a cough at that stage. He was only 76 and was

in great shape physically. He loved to go out and it would not have been a problem for him to go to hospital. The care home called me and said he had symptoms, a bit of a cough and that doctor had assessed him over mobile phone and he would not be taken to hospital. Then I spoke to the GP later that day and said h would not be taken to hospital but would be given morphine if in pain. Later he collapsed on the floor in the bathroom and the care home called the paramedic who established that he had no injury and put him back to bed and told the carers not to call them back for any Covid-related symptoms because they would not return. He died a week later. He was never tested. No doctor ever came to the care home. The GP assessed him over the phone. In an identical situation for someone living at home instead of in a care home, the advice was “go to hospital”. The death certificate says pneumonia and COVID, but pneumonia was never mentioned to us.’

‘A care home manager in Yorkshire told Amnesty International: In March, I tried to get [a resident] into hospital—the ambulance had employed a doctor to do triage but they said, “Well he’s end of life anyway so we’re not going to send an ambulance” ... Under normal circumstances he would have gone to hospital ... I think he was entitled to be admitted to hospital. These are individuals who have contributed to society all their lives and were denied the respect and dignity that you would give to a 42-year-old; they were [considered] expendable.’

The CQC felt it necessary to issue a statement in August 2020 addressing the issue of inappropriate DNR’s being placed on care home residents without informing the resident or their family.

‘It is vitally important that older and disabled people living in care homes and in the community can access hospital care and treatment for COVID-19 and other conditions when they need it during the pandemic ... Providers should always work to prevent avoidable harm or death for all those they care for. Protocols, guidelines and triage systems should be based on equality of access to care and treatment. If they are based on assumptions that some groups are less entitled to care and treatment than others, this would be discriminatory. It would also potentially breach human rights, including the right to life, even if there were concerns that hospital or critical care capacity may be reached.’

That statement was issued because the [CQC found that 34% of people](#) working in health and social care were pressured into placing ‘do not attempt cardiopulmonary resuscitation’ (DNACPR) orders on Covid patients who suffered from disabilities and learning difficulties, without involving the patient or their families in the decision.

The evidence is all there to see in the public domain and thankfully, a team of people are gathering said evidence and analysing it, so that justice can be sought for the people that have had their lives ended prematurely via the use of inappropriate DNR orders, used as permission to start end of life treatment which included a drug called Midazolam. A drug that is associated with respiratory depression and respiratory arrest, the exact same symptoms of complications due to the alleged Covid-19 disease, especially when used for sedation in noncritical care settings.

A drug which was ordered by UK authorities in March 2020 at a quantity to cover a usual two year supply. A two year supply that seems to have been depleted by October 2020 according to NHS documents –

Shortage due to increased demand	Page
Clenil inhaler (and Soprobe)	2
End of Life medicines (general advice)	3
- Levomopromazine injection and tablets	3
- Glycopyrronium injection	3
- Haloperidol injection and oral meds	3
- Midazolam injection	3
- Morphine Injection	3

[Source](#)

But once replenished the stocks were again depleted by the beginning of February 2021 according to official NHS documents –

8th February 2021		Midazolam	10mg/2ml soln for inj in amp	There are limited stocks of midazolam 10mg/2ml solution for injection ampoules available until April 2021. Primary care should: • continue to order midazolam 10mg/2ml solution for injection ampoules in line with current demand for palliative care and are advised not to over order during this time • when prescribing midazolam 10mg/2ml as an anticipatory medication, review the amount required in order to balance timely access with the impact of increased demand on the medicines supply chain in light of the limited stock.
-------------------------	--	-----------	---------------------------------	---

[Source](#)

It was decided in 2013 after a review that the '[Liverpool Care Pathway](#)' was to be abolished. The Liverpool Care Pathway (LCP) was a scheme that we're told intended to improve the quality of care in the final hours or days of a patient's life. It's alleged aim was to ensure a peaceful and comfortable death. The LCP was a guide to doctors, nurses and other health workers looking after someone who was dying on issues such as the appropriate time to remove tubes providing food and fluid, or when to stop medication.

The reason it was decided it should be abolished is that the review found hospital staff wrongly interpreted its guidance for care of the dying, leading to stories of patients who were drugged and deprived of fluids in their last weeks of life.

The government-commissioned review, headed by Lady Neuberger, found that poor training and a lack of compassion on the part of nursing staff was to blame. Harrowing stories from families revealed they had **not** been told their loved one was expected to die and, in some cases, were shouted at by nurses for attempting to give them a drink of water. Nursing staff had wrongly thought, under the LCP guidance, that giving fluids was wrong.

The review made 44 recommendations, including the phasing out of the LCP over six to twelve months as individual care plans for the dying were brought in. It stated that only senior clinicians must make the decision to give end-of-life care, along with the healthcare team, and that no decision must be taken out of hours unless there is a very good reason.

The evidence suggests that the Liverpool Care Pathway returned with a vengeance in April 2020 under the direction of the Health Secretary Matt Hancock, Government Advisors and NHS Chiefs, and it looks as if it was used to manipulate you into giving up over one year of your life under the pretence that you were staying at home, to protect the NHS and save lives. But the evidence suggests that in reality you were ordered to stay at home, to protect the NHS, so that they could prematurely end the lives of the elderly and vulnerable and tell you that they were Covid deaths.

Midazolam. It should be the word that is on everyone's lips. We're sure it will be now.

<https://dailyexpose.co.uk/2021/08/29/midazolam-was-used-to-prematurely-end-the-lives-of-thousands-who-you-were-told-had-died-of-covid-19/>

<https://tapnewswire.com/2021/08/midazolam-was-used-to-prematurely-end-the-lives-of-thousands-who-you-were-told-had-died-of-covid-19-and-we-can-prove-it-heres-the-evidence/>

Before It's News, Sunday, August 29, 2021 5:41

<https://beforeitsnews.com/eu/2021/08/midazolam-was-used-to-prematurely-end-the-lives-of-thousands-who-you-were-told-had-died-of-covid-19-and-we-can-prove-it-heres-the-evidence-2676205.html>

Meer informatie

<https://robscholtemuseum.nl/?s=The+Tap>

<https://robscholtemuseum.nl/?s=David+Icke>

<https://robscholtemuseum.nl/?s=Midazolam>

<https://robscholtemuseum.nl/?s=Covid>

<https://robscholtemuseum.nl/?s=Genocide>